


GEOLOGIC MAP OF THE WILLIAMSBURG 30- x 60-MINUTE QUADRANGLE, VIRGINIA

C.R. BERQUIST, JR.
2013



DESCRIPTION OF MAP UNITS

- Fill** - Undifferentiated material used for fill, modified or disturbed material.
- Marsh sediment** - Organic-rich silt with fine sand and clay interbeds.
- Swamp sediment** - Post and organic-rich sand, silt, and clay.
- Allochthon** - Pebbly sand and sand; silt; interbeds of silt and fine sand with organic matter content.
- Beach and dune** - Fine- to coarse-grained quartz sand, poorly to well-sorted; may include rip-rap adjacent to water.
- Shelly organic mud** - Organic mud with abundant shell material; well-sorted shells or masses of oysters, fossil Rongia, and other shells in contact with one another surrounded by mud; may contain areas of mostly undisturbed silt in place.
- Oysters** - Live oysters and oyster shell; substrate may be mud or sand; shells are typically in contact with one another and forming mounds or reefs; may include some fossil shell.
- Peat** - Decomposed plant remains commonly mixed with sand or mud; dark brown; may include roots of inundated marsh and swamp plants.
- Organic mud** - Organic-rich clay and silt; black, dark gray, or dark brown; consistency ranges from fluid or soupy to somewhat firm.
- Sand** - Fine- to coarse-grained sand; may include granules, small pebbles, and organic material; gray to channel bottom or brown adjacent to cliffs, beaches, and active erosion; along shorelines, it transitions to adjacent mud channels over tens of feet by becoming gradually more muddy and less sandy; contacts that shift with time are shown with a zig-zag pattern.
- Silt and gravel** - Fine- to coarse-grained sand, granules, pebbles, and cobbles; commonly flows typically found adjacent to cliffs or exposed Pleistocene-age deposits.
- Tabb Alteration, undivided** - Clayey fine- to coarse-grained silt.
- Tabb Alteration, Sedgfield Member** - Pebbly sand, and clayey silt.
- Tabb Alteration, Lynnhaven Member** - Silt, fine sand, and clay.
- Tabb Alteration, Lynnhaven and Pappan Member, undivided** - Fine to coarse sand, clayey sand, and silt.
- Tabb Alteration, Pappan Member** - Sand, silt, and clay.
- Easing Green Alteration** - Slightly muddy fine- to medium-grained sand that grades downward to coarse-grained sand, granules and pebbles.
- Shirley Alteration** - Pebble to boulder sand, sand, silt, and clay; scattered post and organic silt beds.
- Chickahominy Alteration** - Pebbly sand, sand, silt, and clay.
- Charles City Alteration** - Pebble to boulder sand, interbedded coarse-bedded sand and silt, and silt and clay.
- Windward Alteration** - Clayey, fine- to coarse-grained sand overlying fine- to coarse-grained sand with pebbles.
- Moorings Unit of Oaks and Coch (1973)** - Massively bedded sand or clay.
- Bacon Castle Formation** - Interbedded sand, silt, and clay; massively bedded and lenticular to flow-bedded basal pebbly to coarse sand contact.
- Chesterbrook Group, Yorktown and Eastern Formations, undivided** - Interbedded shelly fine-grained sand, clayey fine-grained sand and silt; fine to very fine-grained sand. Moore House member: Clayey silt with very fine grained sand grading downward to mud; fine- to coarse-grained sand with granules; glauconitic common.

- Map Symbols**
 - Stratigraphic Contact: approximately located
 - Normal Fault Contact: not exposed at surface; U = upthrown block, D = downthrown block
 - Chlorine Bay Impact Structure
 - Cliff: Marginal of contact center (Henson, Powers, and Goss, 2005)
- INTRODUCTION**

Mapping of the Williamsburg 30- x 60-minute quadrangle began in 1959, in response to rapid growth along the I-64 corridor of the York-James Peninsula. This effort incorporated the early detailed mappings by Bick and Coch (1960) and Johnson (1973), with some modification of their work. The regional stratigraphy has been simplified into a consistent nomenclature, particularly within the Pleistocene map units. This work is not complete. Mapping in the Quaternary quadrangle, east of Richmond, in early 2013 confirmed that consistent sedimentary units of late Quaternary age (Cold Harbor Member) are traceable into the western part of this map area where they comprise the lower part of the Bacon Castle Formation.

A few miles to the west, the Providence Fringe Fault, a high-angle normal fault, is mapped roughly south to north from the James River to the Pamunkey River. This fault is likely synsedimentary with the Richmond Member of the Yorktown Formation with a throw of about 130 feet (down to the east). The effect of faulting on the Yorktown-Eastern contact is evident in the cross-section. Drilling in August 2013 verified westward dipping contacts earlier observed by Johnson (2007), and delineated a normal fault block (see cross-section) related to deformation in the vicinity of the Chesapeake Bay Impact Crater. This deformational event is likely synsedimentary to the Moore House Member. Similar faulting was found in eastern Gloucester County. Additional radial and linear normal faults are likely to be found in the highly uplifted part of the lower York-James Peninsula.

Environmental hazards within the map area include acid forming soils, sinkholes, shoreline erosion, and inundation caused by rising sea level. Acid forming soils result from excavation and exposure of the Eastern Formation. This formation has been mapped in some detail from the Chesapeake Bay to Richmond. Numerous effects of sea level rise, including coastal erosion, salt water intrusion, highway salts, drain clogs, and guardrails, dissolution and staining of concrete, and the inability for re-vegetation without salinity. Shoreline erosion is common within the map area. Global sea level rise coupled with local subsidence will exacerbate this problem.

Mapping bottom sediments in rivers is a recent addition to the Williamsburg quadrangle project. This effort has increased the value of the geologic map in that aquatic habitats have been defined and submerged economic resources have been delineated. Rehabilitation of the American shrewmole relies on identification and protection of hard bottom areas with sparse sediments that serve as spawning grounds. Bottom areas mapped as shelly organic-rich mud are being dredged in the James River as a source for oyster shell to be used in reestablishing depleted oyster reefs elsewhere. These artificial reefs in higher salinity water will provide a new habitat for oyster spat to recolonize and rebuild productive oyster reefs.
- EXPANDED EXPLANATION**

CENOZOIC TO PLEISTOCENE FORMATIONS

Lower Chesapeake Group, Eastern Formation

The Chesapeake Group was defined by Dall and Harris (1963) to include all formations of Miocene age from the Chesapeake Bay to Florida. Ward and Blackwelder (1980) defined the Eastern Formation and its two members, the Clement and Coburn Bay, from exposures on the James River in Surry County, Virginia. Chesapeake (then in the Lower Chesapeake Group) from 1980 to 1981. The Marys Formation was mapped locally to unconformably underlie the Yorktown Formation (Bick and Coch, 1960). The Saint Marys Formation was more recently recognized as the Eastern. For this map, both the Upper Chesapeake Group sediments have been consolidated into the Chesapeake Group (T). We chose not to map the Eastern for the following reasons: 1) At 1:100,000-scale, it was not feasible to show the thin (centimeter to 21) Lower Yorktown (Bathurst and Sunkens Meadow members) shelly sand and lithologically similar to Upper Eastern (Coburn Bay) shelly sands; 2) 3) dissolution of shell materials very common in outcrop and borings; this removal of guide fossils (fossils) impedes interpretation and assignment to a formation most difficult.

Consistent lithologic characteristics have been observed in Chesapeake Group sediments through recent detailed mapping in the York-James Peninsula from Richmond to the Chesapeake Bay. This work includes almost 1,000 auger borings and supports the existing nomenclature with the following lithologic observations: 1) the Coburn Bay sediments, in the absence of fossil material, are composed of slightly muddy to muddy, dark-brown-gray, very fine- to fine-grained sand that commonly contains fine-grained sand-filled burrows, pyrite, iron-sulfide, and 1-5% heavy minerals. This sediment contains, with subtle difference to the overlying fine- to medium-grained Yorktown sands; 2) the Clement Bay sediments are somewhat shelly, dark greenish-gray, very fine-grained mud (compacted and sandy), that commonly contains dark gray fine-grained sand-filled burrows, and some pyrite. Acid-forming soils commonly form on exposure of the Eastern sediments to the atmosphere. The age of the Eastern is Late Miocene. The thickness of this formation above sea level ranges from thin to about 30 feet in the western part of the map area.

Upper Chesapeake Group, Yorktown Formation

The Yorktown Formation was named by Clark and Miller (1966) for exposures on the York River cliffs at Yorktown, York County, Virginia. After numerous modifications and age determinations, Ward and Blackwelder (1980) assigned the Yorktown to the Upper Chesapeake Group and defined four members. Many outcrops and most borings encounter little if any shell material. In the absence of fossil evidence to the contrary, keypoint current nomenclature seems appropriate. The following lithologic characteristics were observed while mapping Richmond and the Chesapeake Bay: 1) the Sunkens Meadow and Richmond Members are shelly, light to bluish-gray, fine-grained mud and silt with 1-2% heavy minerals and glauconite; 2) the overlying Morgans Beach Member is a bluish gray, very fine-grained clayey silt that commonly contains thin beds of disarticulated mollusks. In the Yorktown area, 1-inch to 1-foot-thick beds of shell hash occur sporadically.

The Moore House Member was named for exposures on the York River at Moore House, Colonial National Historical Park. Much of the upper part is covered, but borings have found that this member is quite extensive across the map area. Some sediments formerly mapped as "unsorted" Sundry Formation have been assigned to the Moore House. The Sundry Formation was informally named by Moore (1956) for sediments that overlie the Yorktown Formation and underlie the Bacon Castle Formation. Coch (1962) formalized the name; its use was resumed in subsequent reports (Coch, 1968; Bick and Coch, 1969; Oaks and Coch, 1973). Johnson and Bergquist (1987) first suggested that the Sundry be abandoned. Recent borings at the type and reference sections confirmed that sandy muds assigned to the Sundry are Morgans Beach sediments and as such, abandonment of the name is required.

A thin (less than 18-foot-thick) bed widespread across the map area is represented by "wavy" lines. This bed is the base of the Sundry and is the base of the Yorktown Formation.

The Sundry is a fine-grained, clayey mud to silt, and is the base of the Yorktown Formation and above the Yorktown Formation can be traced from Roanoke (see Providence Fringe Fault) to Yorktown and from Gloucester to Dendron. It is distinct and not to be confused with weathered and non-fossiliferous Sunkens Meadow and Richmond sediments, the Morgans Beach Member of the Yorktown, or the Clement Member Member of the Eastern Formation. This layer was traced to borings eastward to the Moore House type section. The shell hash is a sign of a fossiliferous, but lithologically equivalent and when deeply weathered is the source for much of the clay found locally.

In addition to fossiliferous mud, the Moore House is composed of a sandy clayey-silt mud that is yellowish brown to strong brown in color. The sand fraction coarsens downward from very fine-grained sand to granules. This sequence is typically 7 to 10 feet thick. Coarse sand and granules are commonly very well-sorted. Glauconite is common in the lower part of the sandy mud. Wavy bedding is rarely preserved as extensive laminations has been interpreted in the past. In three places, drilling has encountered a sequence of wavy to yellowish gray, fine- to coarse-grained sand with gray clay layers below the typical sandy mud (and above shelly Richmond/Sunkens Meadow sediments). Some weathered and cemented shell fragments are rarely found.

The thickness of the Yorktown ranges from absent (where removed during sea level lowstands and subsequent deposition of estuarine terraces) to approximately 130 feet. This abnormal thickness occurs east of the north-south trending Providence Fringe Fault, a few miles west of the map area. The age of the Yorktown was previously estimated to be Early Pliocene. Recently, however, Chesapeake and Oaks from an exposure at Tatum Neck (Buckhimer) yield preliminary stratigraphic ages of 2.5 to 0.6 Ma (Paul Hart, personal communication).

Bacon Castle Formation

The Bacon Castle Formation is perhaps the most widespread surficial map unit in the Virginia Coastal Plain. It was informally named by Coch (1965, 1968) where he defined the name for the community of Bacon Castle in Surry County. His type section for wet facies is exposure in road cuts in Clement and Surry quadrangles, also in Surry County. More recently, Ramsey (1988) provided a comprehensive study of the Bacon Castle and defined two members, the Varina Cove and Barhamville. Although this map does not discriminate between these two members, Ramsey's nomenclature redefinition is most reasonable. The Varina Cove member is composed of reddish brown, cross-bedded sand and gravel that grades upward into mud and muddy sand. Lenticular to flaser bedding is common and basal clasts usually contain cobbles. The Barhamville member is composed of variable amounts of sand and mud. Coburn ranges from red and brown to light gray. Lenticular to flaser bedding is also common. Within the map area, thickness ranges upward to about 80 feet. The Bacon Castle is thought to be Late Pliocene in age.

Moorings Unit

The informal "Moorings unit" was defined by Oaks and Coch (1973) and carried forward in the mapping of Moore and others (1980) and the more recent geologic map of Virginia (1983). It originally represented the central and western facies of the Elberton formation (Coch, 1965) and that equivalent in the currently used Williamsburg quadrangle. The Moorings unit is a fine-grained sand and clay (Coch and Coch (1973) provide a description of a core in Runnymede quadrangle that they use to define the Moorings (Section E, Appendix I). It is described as 24 feet of fine-grained sand with some silt. The contact with the underlying Bacon Castle is sharp. Mapping for this project supports their definition and adds two additional borings as reference sections, 2008273 in Surry and 20100343 in Elberton, Dandridge quadrangle. Similar contacts were found at about 105 feet in these borings. The western extent of this Moorings sediment is very difficult to discern from underlying Bacon Castle sediments; both are mud or muddy sands. Note, however, the distinct difference in the degree of dissection above and below 11.5 feet westward of the Surry Survey on 7.5-minute topographic maps. Land surface above 11.5 feet is more dissected than land (less than) 11.5 feet. Possibly there are very thin, eroded, and reflect Moorings mud and muddy sand beneath this line.

Windward Formation

The Windward was named by Coch (1960) for a sand, silt, and clay sequence in a boring near the town of Windward, Isle of Wight County. It has commonly been mapped as a sandy mud grading downward to muddy coarse sand and gravel. This was noted years ago. The Elberton (Windward) is a massive unit 10 feet to 20 feet thick that exhibits no sedimentary structure. It is locally sand, but contains much silt and clay locally. On the whole, the Elberton (Windward) resembles a completely homogenized mixture of the various facies of the Bacon Castle. In some places the basal contact is distinct, in others some gradual gradation (Department of Geology, 1967). A more accurate description of Windward sediments has proven to be problematic in the map area. For example, the Berryman Sand (Surry Quadrangle) shows massive (dissected) gray and light gray mottled muds that grade downward to fine- to coarse-grained sand over a 15-foot interval. These basal sands continue westward and coincide with sand in the lower part of the Bacon Castle Formation. In the map area, the Windward appears to be mostly reworked Bacon Castle sediments that underlies a prominent morphologic setting. The Windward lies below the Surry Survey on 7.5 and 90 feet in elevation and below a scarp rise of 55 feet. Given the problematic lower sands that are likely Bacon Castle sediments, thickness ranges upward to about 15 feet in the map area.

Charles City Alteration

The Charles City was named for a thick sequence of gray to light brown clayey silt, sand, pebbles and cobbles exposed adjacent to the James River in Charles City County, Virginia (Johnson and Bergquist, 1989). These sediments typically lay below facies that range from 60 to 75 feet in elevation and below scarp where the toe is between 75 and 80 feet. Thickness ranges from 60 to 75 feet to a further edge. In the Yorktown area, gray, slightly muddy fine- to medium-grained sands with flaser and lenticular bedding are common. Sediments of the Charles City follow the typical marine transgressive sequence in estuaries by grading upwards from coarse pebbly sand to lenticular-bedded clay-silt. In conjunction with morphologic setting, lagged laminae sands are somewhat correlative to the Charles City; single fine sandy loam soils are also commonly found (USDA 2010a and b). The age of the Charles City is uncertain, but older than the Chickahominy.

Chickahominy Alteration

The Chickahominy was named for an upward-fining sequence of sediments in Isle of Wight County, Virginia by Johnson and Bergquist (1989). Pebble or coarse-grained sand typically grades upward to clayey sand or sandy clay. Organic-rich mud and muddy sand may also be encountered having been deposited in marsh and swamp environments. Sediments are found below facies ranging from 50 to 55 feet in elevation and below scarp where the toe is approximately 58 feet. The age is uncertain, but older than the Shirley (1400,000). It is possible that these sediments relate to a small highland around 500,000 years ago, marine oxygen isotope stage 25.

Shirley Alteration

The Shirley was named for deposits exposed in borrow pits at Shirley Plantation, Charles City County, Virginia (Johnson and Bergquist, 1989). This is composed of a basal gravelly sand grading upward into a fine- to coarse-grained sand and silt, overlain by clayey-silt (mud) or muddy sand. In the Yorktown area at boring location 3428B-1, the Shirley is a light gray silty clay grading downward to pebbly mud, sand, and silt. This is a typical marine transgressive sequence. Consistency of the soils, in part, include Beebea and Chickahominy silty loams. Shirley sediments were deposited during a marine transgression about 400,000 years ago (marine oxygen isotope stage 11). In estuaries, it conforms to the typical upward-fining sequence, but in more coastal areas such as Gloucester Point, it is mostly composed of sand. Shirley sediments underlie facies between about 35 to 40 feet in elevation and below a scarp rise of about 40 feet. Its thickness ranges from a further edge to over 60 feet.

Easing Green Alteration

The Easing Green Alteration was named for sediments that lie beneath a flat at a 30 feet elevation and are bounded by a prominent scarp rise at approximately 38 feet, adjacent to the Pamunkey River, Tazewell 7.5-minute quadrangle west of this map area (Gillmer and Bergquist, in progress). This, slightly muddy fine- to medium-grained sand grades downward to coarse-grained sand, granules and pebbles with about 2% heavy minerals including phosphate locally. The Easing Green sediments are typically sandy throughout the map area. Thickness ranges up to 15 feet but can be much thicker in filled paleochannels in the York and James Rivers.

Recent Lockwood (personal communication) estimated an age of about 125,000 to 100,000 based on amino acid work on Mercuraria and Ostraea in correlative deposits adjacent to the Pamunkey River. Hancock, Gillmer, and Bergquist (in preparation) found a similar cosmogenic age on sand at Easing Green. This age correlates to marine oxygen isotope stage 7a.

A local sea level high-stand at just below 40 feet in elevation may not everywhere have a prominent shoreline (scarp) in guide the delineation of sediment formation contacts. This is a particular area problem to resolve in urbanized settings such as in the City of Newport News. However, soil maps show where contact changes in soil types above and below 35-foot elevation. In addition, accurate ridge defined by the 30-foot contact are found east of Tabb, on the Newport News North 7.5-minute quadrangle (Stage fine sandy loam). These are and are likely older than Tabb Sedgfield sediments. Mapping for this project considered the distribution of Stage, Beebea, Coburn and Alvarado soils (USDA 2010, 2011 and d) in conjunction with the morphologic setting to support the Easing Green in the Newport News area.

On the Hog Island peninsula, there is a clear demarcation of the Easing Green based on morphology and age. In Surry County, Newland silt loam and Alvarado silt loam commonly form on Osg sediments (USDA 2007), in Isle of Wight, Powells silt loam commonly forms on Osg sediments (USDA 2010a); in James City County, Powells and Chickahominy silt loam are common on Osg sediments (USDA 2010a).

Tabb Alteration

The Tabb Formation was named by Johnson (1970) and was subdivided into three members. They are the Sedgfield, Lynnhaven and Pappan Members in descending age and surface elevations. The members are thought to correlate with marine oxygen isotope stages 5c, 5a, and 1, respectively. In its type section in Newport News, the Sedgfield is a fossiliferous, gray to yellowish-brown, cobbly clayey, medium- to coarse-grained sand grading upward to clay or clayey sand, and is about 11 feet thick. Fossils include Mercuraria mercuraria and Crossostraea virginica. In the Chickahominy River, the Sedgfield is commonly sandy and typically contains up to 2% heavy minerals. The Sedgfield is found in thin between 20 to 60 feet in elevation and below a scarp rise of about 10 feet.

The Lynnhaven was named for exposures in a drainage ditch at Butler Farm in Hampton. It is composed of a basal gravelly sand grading upward to bluish-gray sandy clay or clayey sand. In the Chickahominy River, the Lynnhaven is commonly sandy, and typically contains up to 2% heavy minerals. On the Eastern Shore, the Lynnhaven is slightly muddy fine- to medium-grained sand grading downward to medium- to coarse-grained sand, with granules and pebbles. Thin, it correlates with the Newsomford Formation, Oceanbuck Member of Moxon (1985). The Lynnhaven sediments underlie facies between about 60 to 105 feet in elevation and below a scarp rise of about 10 feet.

The Pappan is the Yorktown in the City of Portsmouth, Hampton 7.5-minute quadrangle (Johnson, 1970). On this map, it is restricted to areas with ridge and swale morphology at elevations below 10 feet and to some low-lying nearby areas. On Plum Point Island, it is dark gray silt grading downward to a light gray to reddish brown medium- to coarse-grained sand with pebbles and cobbles and is about 15 feet thick. Heavy minerals comprise 3% of the formation. On Malberry Island, the ridges are capped with fine sandy clay. On Jamestown Island, the Pappan appears to be a sandy clay or clayey sand, grading into the underlying Lynnhaven sediments.

BOTTOM SEDIMENT SAMPLING

Bottom sediment mapping for this project began in 2007 in the Chickahominy River and expanded to the James River within the Surry 7.5-minute quadrangle. This new mapping, shown on the January 2013 version of the Williamsburg sheet covered the York River upstream from the Coleman Bridge, the James River upstream of Malberry Point, the Chickahominy River downstream from Walters dam, and the navigable part of Disappointment Creek. A map from Nichols (1978) was used for the lower James River; a map adapted from Millard (1954) was used for Malberry Bay. Sediment distribution for the rest of the Bay was adapted from Byrne, Hobbs and Caron (1982); the 4-ft contour line separating mud from sand was taken from their Figure 28. Areas of no data are represented by "wavy" lines.

A hand-drawn map of the Bay was taken from their Figure 28. Areas of no data are represented by "wavy" lines. Data from Haven, Whitcomb, and Kennedy (1981) were not included in this version of the map, but might be incorporated in the future along with additional ideation and sampling.

The survey involved making multiple transects across the rivers, stopping at various intervals and recording the station location with GPS. At each station, a grab sample of the bottom sediment was collected with small "diamond" or PVC pipe devices. Transportation at bottom ranged up to about 6 inches. Map units evolved naturally over time with multiple encounters of similar bottom sediments. Contacts between observed materials were marked on 7.5-minute topographic maps in the field and later modified in a GIS setting aided by plotted station locations. In 2009 we began using a Humminbird 119T side-imaging sonar on loan from VMRC/VIMS and later replaced with our own 1190C. The sonar imagery was used in conjunction with sediment samples to make an interpreted map. In addition to samples, bottom material was interpreted by dragging samples (or a chain) in conjunction with observing different imagery patterns. At each station, a "rougher" of the imagery was taken; water depth, bearing, course direction, time were also recorded by the Humminbird. Our tracklines evolved to a rigid pattern of approaching shorelines (typically slightly bottom), then heading in deeper water where black muds typically would be encountered. Linear trends of sediment types were extrapolated between traverses. As confidence in interpretation of the imagery grew, fewer grab samples were collected.

Because this geologic map is regional in scope, some bottom sediment details have been omitted and some areas were not traversed in our survey. A more detailed survey might include additional map units such as brown sand, gray sand, sand and shell, significant man-made objects, and discriminate between different shell types (live oyster, oyster shell, Rangia both live and shell only).

The bottom sediments are continuously moving. For example, a brown sandy shell was found at the south side of the mouth of Queen Creek several years ago. Recent sampling found covering this sand in mud places. Erosion holes of sand marginal to channels gradually grade into mud at up- and down-stream ends; contacts with central channel mud are typically abrupt. Sand waves up to 1 meter in height are common between Hog Island and Jamestown Island and indicate active sand transport, some of which appears in fill in the dredged Goose Hill Channel. This regional map is a snapshot in time of generalized bottom sediment type; contacts will change with time.

ACKNOWLEDGEMENTS

This map was prepared in cooperation with the United States Geological Survey under the National Cooperative Geologic Mapping Program - STATEMAP Component 20120203 Agreement G11AC028. Appreciation is extended to Jack Travelstead of the Virginia Marine Resources Commission for providing a Humminbird side-scan sonar to use for bottom sediment mapping. David Staaboe and Kerry Augusta of Virginia Institute of Marine Science provided technical assistance on use of the side-scan sonar. Any Gillmer, Aaron Cross, Pete Bergquist, Lee Briscoe, Carr Romanach, Dennis Feeley, and Hannah Shepherd provided assistance in the field.

REFERENCES

Bick, K.F., and Coch, N.K., 1969. Geology of the Williamsburg, Hag Island, and Bacon Castle Quadrangles, Virginia. Virginia Division of Mineral Resources, Report of Investigations 18, 28 p.

Byrne, R.L., Hobbs, C.H., III, and Caron, M.J., 1982. Baseline sediment data to determine distribution, physical properties, sedimentation budgets and rates in the Virginia portion of the Chesapeake Bay. Virginia Institute of Marine Science, College of William and Mary, Gloucester Point, Virginia, 152 p.

Clark, W.B., and Miller, R.L., 1906. Clay deposits of the Virginia coastal plain. Virginia Geological Survey Bulletin 2, p. 1.

Coch, N.K., 1965. Post-Miocene stratigraphy and morphology. In: Coastal Plain, southeastern Virginia. Office of Naval Geography, Technical Report of Report 9, 97 p.

Coch, N.K., 1968. Geology of the Beane, Church, Smithfield, Windsor, and Chickahominy quadrangles, Virginia. Virginia Division of Mineral Resources, Report of Investigations 17, 39 p.

Dall, W.L., and Harris, G.D., 1962. Correlation maps. Virginia Geological Survey Bulletin 84, 349 p.

Department of Geology, College of William and Mary, 1967. Geology of the Middle Coastal Plain, Virginia. National Association of Geology Teachers Eastern Section - Spring Meeting, 3 p.

Gillmer, A.K., and Bergquist, C.R., Jr. in progress. Geology of the Tazewell 7.5-minute quadrangle, Virginia.

Haven, D.S., Whitcomb, J.P., and Kendall, P.L., 1981. The present and potential productivity of the Bay of the Chesapeake in Virginia. Volume 2. Virginia Institute of Marine Science, Special report in applied marine science and ocean engineering No. 243.

Horton, L.W., Jr., Powers, D.S., and Goss, C.S., eds., 2005. Studies of the Chesapeake Bay impact structure - The USGS-NASA Langley corals, Hampton, Virginia, and related corals and geophysical surveys. U.S. Geological Survey Professional Paper 1604-A, 433 p.

Johnson, G.H., 1972. Geology of the Yorktown, Potomac West, and Potomac East Quadrangles, Virginia. Virginia Division of Mineral Resources, Report of Investigations 30, 57 p.

Johnson, G.H., 1976. Geology of the Malberry Island, Newport News, North, and Hampton Quadrangles, Virginia. Virginia Division of Mineral Resources, Report of Investigations 41, 72 p.

Johnson, G.H., and Bergquist, C.R., Jr., 1989. Geology and mineral resources of the Beane and Noyes quadrangles, Virginia. Virginia Division of Mineral Resources Publication 87, 28 p.

Milligan, D.A., 1994. An investigation of the Late Quaternary morphology of Malberry Bay, Virginia and application of a facies model. M.S. Thesis, Virginia Institute of Marine Science, 63 p.

Mixon, R.B., 1985. Stratigraphic and geomorphic framework of uppermost Cenozoic deposits in the Southern Delmarva Peninsula, Virginia and Maryland. U.S. Geological Survey Professional Paper 1067, 63 p.

Mixon, R.B., Powers, D.S., Ward, L., and Andrews, G.W., 1981. Lithostratigraphy and molluscan and diatom biostratigraphy of the Haysville corals, northeastern Virginia Coastal Plain. Chapter A, in: Moxon, R.B., ed., Geology and paleontology of the Haysville corals, Richmond County, northeastern Virginia Coastal Plain. U.S. Geological Survey Professional Paper 1489, 1489 p.

Moore, W.E., 1956. Pleistocene terraces south of the James River, Virginia. Virginia Academy of Geosciences Section Guidebook, May 1956, 16 p.

Nichols, M., 1967. Cruise Guide to the James River Estuary on R/V Langley. Virginia Academy of Science 40th Annual Meeting, map 79.

Nichols, M., 1978. Response and recovery of an estuary following a river flood; Journal of Sediment Res. 47, p. 171-186.

Oaks, R.Q., Jr., and Coch, N.K., 1973. Post-Miocene stratigraphy and morphology, southeastern Virginia. Virginia Division of Mineral Resources Bulletin 32, p. 1-135.

Ramsey, R.W., 1988. Stratigraphic and geomorphic framework of uppermost Cenozoic deposits in the Southern Delmarva Peninsula, Virginia and Maryland. U.S. Geological Survey Professional Paper 1067, 63 p.

US Department of Agriculture, 2007. Soil Survey Geographic Database for Surry County, Virginia. file va073 available at <http://SoilDataMart.nrc.usda.gov>

US Department of Agriculture, 2008. Soil Survey Geographic Database for Mathews County, Virginia. file va115 available at <http://SoilDataMart.nrc.usda.gov>

US Department of Agriculture, 2010a. Soil Survey Geographic Database for Gloucester County, Virginia. file va073 available at <http://SoilDataMart.nrc.usda.gov>

US Department of Agriculture, 2010b. Soil Survey Geographic Database for Isle of Wight County, Virginia. file va093 available at <http://SoilDataMart.nrc.usda.gov>

US Department of Agriculture, 2010c. Soil Survey Geographic Database for James City and York Counties and Williamsburg, Virginia. available at <http://SoilDataMart.nrc.usda.gov>

US Department of Agriculture, 2010d. Soil Survey Geographic Database for Tidewater Cities Area, Virginia. file va115 available at <http://SoilDataMart.nrc.usda.gov>

Ward, L.W., and Blackwelder, B.W., 1980. Stratigraphic revision of upper Miocene and lower Pliocene beds of the Chesapeake Group, middle Atlantic Coastal Plain. U.S. Geological Survey Bulletin 1482-D, p. 61 p.

SUGGESTED REFERENCE

Bergquist, C.R., 2013. Geologic Map of the Williamsburg 30- x 60-minute quadrangle, Virginia. Virginia Division of Geology and Mineral Resources Open-File Report 2013-07, 1:100,000-scale map.



Figure 2. Sinkhole in Colonial National Historical Park battlefield. Dissolution of underlying shells in the Yorktown Formation beneath a void over time, eventually overlying sediments collapse into the void. This is a common occurrence within the map area and in other places in the southern part of the Virginia coastal plain.



Figure 3. Continuous shoreline erosion at York River State Park.



Figure 4. Yellow (silt) coating on outcrop of Eastern Formation, Clement Manor Member from weathering of pyrite results in acid sand with low pH. Fishback Landing, Chickahominy River.




Figure 5. Side-scan image and chart showing transition from shelly organic mud to firm mud, the James River offshore of Kingsmill.




Figure 6. Shell hash - corals, Moore House Member of the Yorktown Formation, forming "Cornwallis Cave" at Yorktown.